

**HURON PERTH HEALTHCARE ALLIANCE
OCCUPATIONAL HEALTH MANUAL**

Healthcare Provider Immunization Status and Surveillance

Appendix A

IMMUNIZATION REQUIREMENTS	
Vaccine or Test	Requirements
COVID-19	Highly recommended.
Hepatitis B	Documentation of Hepatitis B vaccination series (2 or 3 vaccines) and proof of immunity by serology. <ul style="list-style-type: none"> Vaccination with subsequent proof of immunity is highly recommended for Staff who may have exposure to human blood and body fluids. Hep B is not mandatory for volunteers. Individuals refusing the Hepatitis B vaccine must sign a Declination Form.
Influenza	Highly recommended each year. <ul style="list-style-type: none"> It is <u>mandatory</u> to report your Influenza immunization status Annually with the declination form provided in-between the time period (Required October 1 – March 31)
Measles	Laboratory evidence of immunity (titres), OR Documentation of 2 doses with live measles -containing vaccine on or after the 1 st birthday, with doses given at least 4 weeks apart
Meningococcal (Microbiology staff)	It is recommended that Microbiology Technologists who may be routinely exposed to cultures of N. Meningitides receive the following vaccines: <ul style="list-style-type: none"> Quadrivalent Meningococcal A,C,Y,W-135 (Menactra, Menveo, Nimenrix, MenQuadfi,), Re-Vaccination is generally recommended every 5 years if at on-going risk for Men-C-ACYW- 135 Meningococcal vaccine is not routinely recommended for most HCWs.
Mumps	Laboratory evidence of immunity (titres), OR Documentation of 2 doses with mumps -containing vaccine on or after the 1 st birthday, with doses given at least 4 weeks apart
Polio	Recommended completion of initial primary series (3 doses of IPV Polio)
Rubella	Laboratory evidence of immunity (titres), Documentation of 1 dose with live rubella -containing vaccine on or after the 1 st birthday.
TB Skin Test (TST)	You must have documentation of a previous negative 2-step TB skin test. Will not be able to administer TB Skin test if had LIVE Vaccines within the last 30 days <ul style="list-style-type: none"> If you've never been tested, or do not have documentation of a previous negative 2-step, then a 2-step TB test is required If you have documentation of a negative Tb skin test within the past 12 months, no further TB skin testing will be required depending on individual risk assessment For Positive TB reactors, a chest x-ray completed within the past 12 months, and a copy of the x-ray report is required. Please include a copy of previous positive test result. Anyone with a confirmed positive Tb skin test, who has not received counselling or advice concerning Prophylactic treatment, should be referred to Public Health as well as their Primary Care Provider. Chest x-ray will be performed for all new Positive Tb Skin tests
Tetanus/Diphtheria/Pertussis (Tdap) Tetanus/Diphtheria (Td)	Documentation of Td or Tdap vaccination within the last 10 years. <ul style="list-style-type: none"> A onetime adult dose of Tdap (Tetanus/Diphtheria/<u>Pertussis-for whooping cough</u>) is required for all adults if not already received in Adulthood (18 years and older) Tetanus and Diphtheria is recommended every 10 years Recommended completion of initial primary series (at least 3 doses of Td) Tdap given first if >18 years of age
Varicella	Laboratory evidence of immunity (titres), OR Documentation of 2 doses with Varicella-containing vaccinations, given a minimum of 6 weeks apart